

10/556492

Rec'd PCT/PPO 3/25 MAY 9 2005

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



(43) International Publication Date
10 June 2004 (10.06.2004)

PCT

(10) International Publication Number
WO 2004/048369 A1

(51) International Patent Classification⁷: C07D 409/04

301-105 Kwanakseongwon Apt., Buheung-dong, Dongan-gu, 431-054 Anyang-city, Kyungki-do (KR).

(21) International Application Number:
PCT/KR2003/002552

(74) Agent: LEE, Young-Pil; The Chunghwa Building, 1571-18 Seocho-dong, Seocho-gu, Seoul 137-874 (KR).

(22) International Filing Date:
25 November 2003 (25.11.2003)

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: Korean
(26) Publication Language: English
(30) Priority Data:
10-2002-0074119
26 November 2002 (26.11.2002) KR

(84) Designated States (*regional*): ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/048369 A1

(54) Title: METHOD FOR PREPARING OLТИПРАЗ

(57) **Abstract:** Provided is a method for preparing oltipraz. The method includes reacting methyl 2-methyl-3-(pyrazin-2-yl)-3-oxopropionate with phosphorus pentasulfide in the presence of a mixed solvent of toluene and xylene, followed by recrystallization.